



Case Report / Olgu Sunumu

## A very rare renal venous anomaly: Accessory polar left renal vein

*Çok nadir bir renal venöz anomaly: Aksesuar polar sol renal ven*

Ayşe Gül Karadeniz<sup>1</sup>, Hatice Ayça Ata Korkmaz<sup>2</sup>, İsmail Gökhan Çavuşoğlu<sup>1</sup>

*Institution where the research was done:*

Trabzon Ahi Evren Thoracic and Cardiovascular Surgery Training and Research Hospital, Trabzon, Turkey

*Author Affiliations:*

<sup>1</sup>Department of Radiology, University of Health Science,  
Trabzon Ahi Evren Thoracic and Cardiovascular Surgery Training and Research Hospital, Trabzon, Turkey

<sup>2</sup>Department of Radiology, University of Health Science, Trabzon Kanuni Training and Research Hospital, Trabzon, Turkey

### ABSTRACT

A 49-year-old female patient with hypertension was admitted with an intraabdominal murmur and referred for multidetector computed tomography angiography for suspected renal artery stenosis. Her medical history, clinical examination, serum biochemistry, and ultrasound examination findings showed no pathological findings of diabetes mellitus or any other disease. Axial and coronal reformatted multidetector computed tomography angiography revealed a dilated accessory polar renal vein which drained the venous circulation of the left lower pole into the inferior vena cava at midline. In conclusion, multidetector computed tomography angiography plays a progressively substantial role in the assessment of renal vascularity. Vascular surgeons and urologists should be a thorough knowledge of renal vascular variations.

**Keywords:** Accessory renal vein, multidetector computed tomography, polar renal vein, transplant surgery, venous anomaly.

In clinical anatomy, renal venous system has been less studied than its arterial counterpart. Renal vein variations are usually overlooked by radiologists. These variations usually indicate the type and course of the operation, particularly for vascular surgery.

Preoperative description of venous anatomy using multidetector computed tomography (MDCT) angiography is useful to evaluate the arterial anatomy. Recent anatomical studies on renal vascular variations have shown that renal venous anatomical variations are relatively small, compared to arteries.<sup>[1,2]</sup>

### ÖZ

Hipertansif 49 yaşında kadın hasta, abdominal üfürüm ile başvurdu ve renal arter darlığı şüphesi ile çokkesitli bilgisayarlı tomografi anjiyografiye yönlendirildi. Öykü, klinik muayene, serum biyokimya ve ultrason tetkiki bulguları diabetes mellitus ya da başka bir hastalık açısından bulgular göstermiyordu. Aksiyal ve koronal çokkesitli bilgisayarlı tomografi anjiyografi reformat görüntülerde, sol böbrek alt polün venöz dolaşımını orta hatta inferior vena kavaya drene eden dilate aksesuar polar renal ven izlendi. Sonuç olarak, çokkesitli bilgisayarlı tomografi anjiyografi böbrek vaskülaritesini değerlendirmede giderek artan bir role sahiptir. Vasküler cerrahların ve ürologların böbreğin vasküler varyasyonları hakkında detaylı bilgi sahibi olması gereklidir.

**Anahtar sözcükler:** Aksesuar renal ven, çok kesitli bilgisayarlı tomografi, polar renal ven, transplant ameliyatı, venöz anomali.

To the best of our knowledge, renal venous anatomic anomaly with an accessory renal vein has only been reported in one report in the literature.<sup>[3]</sup> Although the definition of accessory renal artery is widely used, the definition of accessory renal vein is little known by radiologists and clinicians. However, recognition of this variation is particularly important in surgical procedures, such as vascular and general surgery.

Herein, we describe the MDCT angiographic features of a very rare venous anatomic anomaly of the left kidney with an accessory renal vein from left renal

Received: November 26, 2018 Accepted: December 22, 2018 Published online: April 24, 2019

**Correspondence:** Hatice Ayça Ata Korkmaz, MD. SBÜ Trabzon Kanuni Eğitim ve Araştırma Hastanesi Radyoloji Kliniği, 61250 Ortahisar, Trabzon, Turkey.  
Tel: +90 538 - 848 22 07 e-mail: dr.h.aycaatakorkmaz@gmail.com


**Cite this article as:**

Gül Karadeniz A, Ata Korkmaz HA, Çavuşoğlu İG. A very rare renal venous anomaly: Accessory polar left renal vein. Turk Gogus Kalp Dama 2019;27(2):241-244

inferior pole which was related to the lower pole of the kidney rather than the renal hilus. We believe that our report would provide a significant contribution to the current literature.

### CASE REPORT


A 49-year-old female patient with hypertension was admitted with an intraabdominal murmur and referred for MDCT angiography for suspected renal artery stenosis. Her medical history and clinical examination revealed no pathological findings of diabetes mellitus or any other disease. Complete blood count and all serum biochemistry were within normal limits. Ultrasound showed no structural anatomic abnormality of the urinary tract. Axial, coronal, and three-dimensional reformatted examinations demonstrated a dilated accessory polar renal vein which drained the venous circulation of the left lower pole into the inferior vena cava (IVC) at midline (Figures 1, 2, and 3). The drainage site of left renal vein to the IVC was just opposite to the drainage site of right renal vein, but the drainage site of the accessory left renal vein to the IVC was just proximal to bifurcation of the common iliac veins, 8 cm below the drainage site of the left renal vein. The main parent left renal vein was in its normal position. Both kidneys, ventral, and dorsal branches were draining into the IVC directly. No renal vein variation or supernumerary vein was observed in the right kidney. Renal arteries of both kidneys originated from their




**Figure 2.** Coronal reformatted computed tomography image showing an accessory renal vein draining the left lower pole, while left main renal vein was localized in its normal position (large arrow).



**Figure 1.** The arterial phase of the axial computed tomography showing an accessory left renal vein (white arrow), lumbar, and anterior external venous plexus drainage.



**Figure 3.** Three-dimensional volume rendering computed tomography image showing an accessory renal vein draining the left lower pole (white arrows).




**Figure 4.** Illustration of normal renal vascularity.

normal position and no renal artery variation was observed. The widths of ureters were normal and no double collecting system was observed. The hypertension of the patient was considered as essential HT because of the normal renal artery vascular structures in MDCT. After medical treatment, the patient's blood pressure was within normal limits and no additional treatment was required for the incidentally detected venous anomaly.

## DISCUSSION

In clinical anatomy, the renal venous system has been under studied than the arterial system. Renal veins are formed by the confluence of two or three renal parenchymal veins in the renal sinus.<sup>[4]</sup> At the eighth week of embryological development, the symmetrical cardinal veins on both sides form the IVC on the right side. Initially, renal veins are on the ventral and dorsal plane, although they join into a single vessel formation later (Figure 4).<sup>[5]</sup> When these branches fail to join and persist, accessory and additional veins are formed.

Any accessory vessel originating from the kidney and draining independently into the IVC is considered a normal variation and such vessels are termed as additional renal veins, and classified as type 3.<sup>[6,7]</sup> The drainage patterns and variations of the renal veins have been well-documented in previous reports (Figure 5).<sup>[3,7,8-12]</sup> Additional renal veins are more common on the right side than the left side.<sup>[6]</sup> In our case, in contrast to all cases described in the literature, a left-sided accessory polar renal vein was observed which was related to the lower pole of the kidney rather than the renal hilus. Gupta *et al.*<sup>[9]</sup> reported an incidence of 33% for a right-sided additional renal vein and 3.3% for a left-sided additional renal vein.



**Figure 5.** Types of left renal vein anomalies. Type 1 retroaortic left renal vein, type 2 retroaortic left renal vein, type 3 circumaortic left renal vein, and type 4 renal vein joining left common iliac vein.

Renal vein variations, which are usually overlooked by radiologists, are often demonstrated by venography or cadaveric dissections. In our case, the right and left main renal veins drain into the IVC in their normal localizations, whereas the left-sided accessory or additional renal vein drains into the IVC proximally to bifurcation of common iliac veins 8 cm below the drainage site of the left renal vein. Based on our literature search, we were able to find only one publication of anatomic variation of accessory polar vein draining the left pole of the left kidney.<sup>[3]</sup>

Renal vascular variations are clinically silent and, thus, often remain unnoticed. They are usually diagnosed incidentally during surgery or autopsy. Of note, MDCT angiography plays a progressively substantial role in the assessment of the renal vascularity.<sup>[4,10-15]</sup> This technique is a less invasive imaging modality which provides a more rapid, effective, and definitive visualization of many anatomical variations of the kidney vessels. Recognition of the renal venous morphology also affects the type and course of the operation. The importance of familiarity with the renal artery variations has been gradually increasing not only for successful vascular surgeries, but also other types of surgery and radiological examinations. Similarly, recognizing renal venous system anomalies provides comfort for surgeons prior to interventional and surgical operations, such as nephrectomy and vena cava filter placement.

In conclusion, renal vein variations often remain unnoticed, as they are clinically silent. Vascular surgeons and urologists should have a thorough knowledge of renal vascular variations.

## Declaration of conflicting interests

The authors declared no conflicts of interest with respect to the authorship and/or publication of this article.

### Funding

The authors received no financial support for the research and/or authorship of this article.

### REFERENCES

1. Patil AB, Javali TD, Nagaraj HK, Babu SMLP, Nayak A. Laparoscopic donor nephrectomy in unusual venous anatomy - donor and recipient implications. *Int Braz J Urol* 2017;43:671-8.
2. Potaliya P, Sharma S, Kataria D, et al. Multiple renal vascular variations: A case Series. *Int J Anat Var* 2018;11:90-2.
3. Hazirolan T, Öz M, Türkbey B, Karaosmanoglu AD, Canyiğit M. CT angiography of the renal arteries and veins: normal anatomy and variants. *Diagn Interv Radiol* 2011;17:67-73.
4. Türkvatan A, Ozdemir M, Cumhuri T, Olçer T. Multidetector CT angiography of renal vasculature: normal anatomy and variants. *Eur Radiol* 2009;19:236-44.
5. Williams PL, Warwick R, Dyson M, Bannister LH, editors. *Gray's Anatomy*. Edinburgh: Churchill Livingstone; 1995.
6. Satyapal K. The renal veins: a review. *Eur J Anat* 2003;7(Suppl. 1):43-52.
7. Satyapal KS. Classification of the drainage patterns of the renal veins. *J Anat* 1995;186:329-33.
8. Satyapal KS, Kalideen JM, Haffejee AA, Singh B, Robbs JV. Left renal vein variations. *Surg Radiol Anat* 1999;21:77-81.
9. Gupta A, Gupta R, Singal R. Congenital variations of renal veins: embryological background and clinical implications. *J Clin Diagn Res* 2011;5:1140-3.
10. Kumar N, Swamy R, Patil J, Nayak S, Shetty S, Aithal A. Accessory right renal vein draining into inferior vena cava. *OA Case Rep* 2014;3:48.
11. Krishnasamy N, Somayaji S, Koshy S, Rodrigues V. An unusual case of unilateral additional right renal artery and vein. *Int J Anat Var* 2010;3:1.
12. Karau PB. Case Report: Supernumerary right renal vein draining inferior to the main renal vein: A case report and review of the literature. *Anat J Afr* 2013;2:132-4.
13. Monkhouse WS, Khalique A. The adrenal and renal veins of man and their connections with azygos and lumbar veins. *J Anat* 1986;146:105-15.
14. Arévalo Pérez J, Gragera Torres F, Marín Toribio A, Koren Fernández L, Hayoun C, Daimiel Naranjo I. Angio CT assessment of anatomical variants in renal vasculature: its importance in the living donor. *Insights Imaging* 2013;4:199-211.
15. Urban BA, Ratner LE, Fishman EK. Three-dimensional volume-rendered CT angiography of the renal arteries and veins: normal anatomy, variants, and clinical applications. *Radiographics* 2001;21:373-86.