

Physician - Aortic (Abdominal) Pathologies and Surgery/Endovascular Interventions

[MSB-34]

Clinical Outcomes and Applications of Bypass Procedures on Intra-Abdominal Visceral Arteries

Muhammet Fethi Sağlam, Emrah Uğuz

Department of Cardiovascular Surgery, Ankara Yıldırım Beyazıt University, Ankara, Türkiye

Türk Gogus Kalp Dama 2024;32(Suppl 2):MSB-34

Doi: 10.5606/tgkdc.dergisi.2024.msb-34

E-mail: dr.m.fethisaglam@gmail.com

Received: September 12, 2024 - Accepted: September 29, 2024

Objective: This study aimed to evaluate the clinical effectiveness, indications, and patient profiles of bypass procedures performed on intra-abdominal visceral arteries.

Methods: Ten patients who underwent between 2019 and 2024 were included in the study.

Results: In five patients (4 females, 1 male; range, 38 to 64 years), the procedure was part of a hybrid treatment due to aortic pathology. The procedure was performed due to mesenteric ischemia in three patients (2 males, 1 female; range, 50 to 65 years), following left renal artery injury in one female patient (aged 67 years), and due to a pancreatic tumor in another female patient (aged 57 years), which required a bypass to the hepatic artery. During follow-up, no early mortality was observed.

Conclusion: Bypass procedures on intra-abdominal visceral arteries yield successful outcomes in treating arterial stenosis causing mesenteric ischemia, as part of hybrid treatments for aortic pathologies, and in abdominal surgeries requiring revascularization.

Keywords: Abdominal aortic aneurysm, endovascular aneurysm repair, occlusive mesenteric ischemia.

References

1. Gupta PK, Horan SM, Turaga KK, Miller WJ. Chronic Mesenteric Ischemia: Endovascular versus Open Revascularization. *J Endovasc Ther* 2010;17:540-9.
2. Oderich GS, Malgor RD, Ricotta JJ 2nd. Open and endovascular revascularization for chronic mesenteric ischemia: Tabular review of the literature. *Ann Vasc Surg* 2009;23:700-12. doi: 10.1016/j.avsg.2009.03.002.
3. Karita R, Koizumi S, Kubota Y, Ueda H, Ishida K. Endovascular aneurysm repair with mesenteric artery bypass for abdominal aortic aneurysm with occlusion of celiac and superior mesenteric arteries. *J Vasc Surg Cases Innov Tech* 2022;9:100927. doi: 10.1016/j.jvscit.2022.04.017.